



# Quat Tuberculocidal HUSKY® 814

## Q/T TUBERCULOCIDAL SPRAY DISINFECTANT

### SPECIFICATIONS:

Registrations	EPA
Dilution	RTU
pH Level	11.8 – 12.8
Color	Clear
Fragrance	Lemon

Effective against  
**Mycobacterium  
tuberculosis in  
5 MINUTES**  
and **Norovirus in  
30 SECONDS.**



A ready-to-use, quaternary ammonium disinfectant spray cleaner that is ideal for healthcare facilities due to its extensive list of disinfectant claims including Tuberculosis and Norovirus. Inhibits growth of mold and mildew and controls related odors. EPA registered as a Broad Spectrum Disinfectant cleaning product.

### FORMULA FEATURES:

- EPA Approved for Use with Check Mate Electrostatic Cleaning System
- Cleaner, Deodorizer and Mildewstat
- Effective Tuberculocidal Disinfectant on Hard, Non-Porous Surfaces
- Unlike Bleach, Will Not Damage Carpets, Upholstery, Clothing or Metal
- Kills Many Viruses and Bacteria in 1, 3 or 5 Minutes
- Kills Norovirus in 30 Seconds
- EPA Registered to Kill Hepatitis A
- Bloodborne Pathogen Compliant

### RECOMMENDED FOR:

- Hard, non-porous surfaces
- Floors
- Walls
- Counters
- Sinks
- Appliances
- Metal Surfaces
- Stainless Steel Surfaces
- Glazed Porcelain
- Ceramic Tile
- Plastic Surfaces
- Bathrooms
- Shower Stalls and Bathtubs

### DIRECTIONS:

**Disinfection, deodorizing and cleaning:** Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

# Q/T TUBERCULOCIDAL SPRAY DISINFECTANT

## EMERGING PATHOGEN COMPLIANCE GUIDANCE

This product has demonstrated effectiveness against viruses similar to SARS-COV-2 on hard, non-porous surfaces. Therefore, this product can be used against COVID-19 when used in accordance with the directions for use against Canine Parvovirus.

### 1/2 MINUTE CONTACT TIME:

- Norovirus
- Feline Calicivirus
- Rabies Virus

### 1 MINUTE CONTACT TIME:

- HIV-1 [AIDS Virus]
- Hepatitis B
- Hepatitis C
- Bovine Viral Diarrhea Virus
- Influenza Type A / Hong Kong [Influenza]

### 2 MINUTE CONTACT TIME:

- Avian Influenza A (H3N2 & H3N9)
- Human Coronavirus
- SARS associated Coronavirus

### 3 MINUTE CONTACT TIME:

- Pseudomonas aeruginosa
- Salmonella enterica
- Staphylococcus aureus
- Vanomycin Resistant Enterococcus faecalis
- Escherichia coli
- Listeria Monocytogenes
- Yersinia enterocolitica
- Streptococcus pyogenes
- Corynebacterium ammoniagenes
- Human Rotavirus
- Rotavirus
- Paramyxovirus
- Methicillin Resistant Staphylococcus epidermis
- CA - Methicillin Resistant Staphylococcus epidermis
- Methicillin Resistant Staphylococcus epidermis
- Rhinovirus Type 36
- Vancomycin intermediate resistant Staphylococcus aureus

### 10 MINUTE CONTACT TIME:

- Mycobacterium bovis (BCG)
- Canine Parvovirus
- Trichopyton mentagrophytes
- Hepatitis A Virus

This is not a complete list of organisms and names